

Review Article

Open Access

Prevalence and Epidemiology of Diabetes Mellitus Type 3C

Samreen Riaz*

University of the Punjab, Lahore, Pakistan.

Abstract


Type-II diabetes mellitus is a particularly heterogeneous and multifactorial metabolic disorder that is characterized by a combination of insulin insufficiency and insulin resistance. The prevalence of diabetes mellitus has risen exponentially over the last three decades, with resultant increase in morbidity and mortality mainly due to its complications. There is need of Increases awareness of these complications among patients and populations, this facilitates early detection and prevention of these diseases. The earlier diagnosis, timely appropriate treatment may help in avoiding severe, irreversible disability and deformity, which can be done with the tool of patient education and conducting seminar to increase awareness.

Keywords: diabetes mellitus; high blood glucose levels; pancreatitis

Introduction


Diabetes mellitus is one of the most prevalent metabolic disorders in the world, it is characterized by the high blood glucose levels (Hyperglycemia) due to the defects in the action of insulin, secretion of insulin or may be due to both of them [1]. Type I and Type II diabetes are considered as the major types of the diabetes to be known by that time and type II Diabetes is more prevalent (>85%) then that of the type I [2]. Type I Diabetes is considered to be caused by the deficit insulin secretion from the beta cells of

pancreas that actually happens due to the autoimmune devastation of the beta cells. On the other hand, diabetes type II is caused by the two coupled reasons, first one is the resistance to insulin action and second one is the insufficient compensatory insulin secretory response. “Gestational Diabetes mellitus” (GDM), that is also a situation of glucose intolerance, is known to be recognized during the pregnancy. Except of these above-mentioned types of Diabetes, there are some more specific categories of the diabetes mellitus and the Type 3c Diabetes fall in this category [3-5].



Among the people who are suffering from the Diabetes mellitus, approximately 2 percent of them have the type 3c diabetes [6-10]. Type 3c diabetes might happen due to genetic as well as the non-genetic factors that comprises pancreatic removal, pancreatitis, pancreatic ductal adenocarcinoma,

hemochromatosis or cystic fibrosis. One of the most dominating causes of the Type 3c diabetes is chronic pancreatitis. Unfortunately, patients suffering from the type 3c diabetes are not diagnosed properly and timely that is one of the main causes of delay in their required treatment [11-15].



Type 3c Diabetes Mellitus (T3cDM)

Type 3c diabetes mellitus has been designated as the secondary diabetes and usually called as the pancreatogenic diabetes. One can also develop type 3c if he/she has a part or all of the pancreas removed

because of any other damage. Some people who take steroids can also develop this type of diabetes. This is also called as steroid-induced diabetes, and is more common in the group of people which are at high risk of type II diabetes.

Table 1A: Sex and age distribution across groups.

	Groups	n	%	Epidemiology					
				Male	Female	Male %	Female %	Age (Average/y)	Age (SEM/y)
	AP	983	74.80%	525	458	53.40%	46.60%	56.7	0.55
RAP-2	2 nd AP	173	13.20%	111	62	64.20%	35.80%	52.6	1.21
RAP-3	3 rd AP	43	3.30%	25	18	58.10%	41.90%	51.9	2.25
RAP-4	4 th AP	24	1.80%	16	8	66.70%	33.30%	50.7	3.59
RAP-5+	5 th AP	30	2.30%	23	7	76.70%	23.30%	52.57	3.56
	RAP	270	20.50%	175	95	64.80%	35.20%	52.7	0.93
	CP	62	4.70%	45	17	72.6%*	27.40%	55.5	1.84
	Total	1315	100.00%	745	570				

* $p < 0.05$ when comparing AP and CP; $p < 0.05$, AP: Acute Pancreatitis, RAP: Recurrent Acute Pancreatitis, CP: Chronic Pancreatitis, n: number, SEM: Standard Error of the Mean, y: Years

Table 1B: Etiology of acute pancreatitis across study cohorts.

Groups	Leading Etiology					
	AP		RAP		CP	
	n	%	n	%	n	%
Biliary	493	50.2%	63	20.7%	7	11.3%*
Alcohol	191	19.4%	119	39.1%	32	51.6%*
Lipid	48	4.9%	29	9.5%	1	1.6%
Idiopathic	189	19.6%	72	23.8%	18	29.0%

* $p < 0.05$ when comparing AP and CP.

Table 1C: Etiology of recurrent acute pancreatitis.

Groups	Leading Aetiology							
	RAP-2 (n=173)		RAP-3 (n=43)		RAP-4 (n=24)		RAP-5+ (n=30)	
	n	%	n	%	n	%	n	%
Biliary	43	24.9%	4	9.3% ^b	2	8.3% ^c	6	20%
Alcohol	65	37.6%	21	48.8% ^b	9	37.5%	12	40.0%
Lipid	17	9.8%	1	2.3%	3	12.5%	3	10.0%
Idiopathic	41	23.7%	12	27.9%	6	25.0%	5	16.7%

a: $p < 0.05$ when comparing AP and RAP2. b: $p < 0.05$ when comparing AP and RAP3. c: $p < 0.05$ when comparing AP and RAP4. d: $p < 0.05$ when comparing AP and RAP5+.


Table 2: Clinical Characteristics for Common Forms of Diabetes.

	T1DM	T2DM	T3cDM		
Associated With	Autoimmunity	Obesity	Chronic Pancreatitis	Cystic Fibrosis	Pancreatic Resection
Median Age of Onset	2nd decade of life	6th decade of life	5th decade of life	3rd decade of life	Within 5 yrs. of surgery
Pancreatic Insufficiency	No	No	Yes	Yes	Yes
Hepatic Insulin Sensitivity	Normal or decreased	Decreased	Normal or decreased	?	Normal or decreased
Peripheral Insulin Sensitivity	Normal or decreased	Decreased	Normal	?	Normal
Diabetic Ketoacidosis	Yes	No	No	No	No
Hypoglycemia Risk	Increased	Normal	Normal or increased	Normal or increased	Normal or increased
Pancreatic Polypeptide Response	Normal or decreased	Normal or increased	Decreased or Absent	Absent	Absent

Pathogenesis of Type 3c Diabetes mellitus

The pathogenesis of T3cDM is characterized by the declination in the insulin production that might be due to the reduced number of the islet's cells or due to the loss of functionality that is attributed to the


fibrosis or sclerosis [16-20]. Basically, the damage of the pancreatic tissue causes the deficiency of insulin and its mechanism can be explained by the functionality dependent interplay between the acinar cells and islets cells.



Pancreatic Function

Pancreas holds two significant functions in the human body, firstly, it controls the blood glucose levels and secondly, it assists in the digestion of the food. Anatomically, pancreas is divided into the five regions, it is associated with the gall bladder, duodenum and spleen. It has both the exocrine and the endocrine regions but the maximum region of the pancreas owns exocrine function that is responsible for the formation and release of the digestive enzymes into the duodenum. Acinar cells that make up to 85 percent of the pancreatic cells are devoted for

synthesizing the enzymes for carbohydrates, lipids and protein digestion. The main enzyme contributors include amylases, lipases and trypsin and zymogens (proelastase, trypsinogen, chymotrypsinogen procarboxypeptidase). Endocrine secretory tissues residing in the pancreas are called as the islets of langerhans, containing four types of the cells, out of them beta cells (Insulin secreting cells) are in the huge amount. In Type I and Type II Diabetes mellitus, deficient insulin production leads to the hyperglycemia but the digestion of food is unaffected. On the other hand, in Type 3c Diabetes mellitus, digestion of food also gets effected.




Causes of Type3c Diabetes Mellitus

One of the most frequently recognized causes of the Type 3c diabetes mellitus is chronic pancreatitis. The most commonly identified cause of type 3c diabetes is chronic pancreatitis. According to a single-center review, causes of the type 3c diabetes were recorded as: “chronic pancreatitis” (79%), “pancreatic ductal adenocarcinoma” (8%), “haemochromatosis” (7%), “cystic fibrosis” (4%), and previous pancreatectomy (2%), results are also depicted in the Pie Chart [21-25].

Chronic Pancreatitis and Type 3c Diabetes mellitus

Progressive inflammation in the pancreatic tissue results in the chronic pancreatitis. Genetic and environmental factors both are considered as its

cause. It is characterized by the gradual devastation of the pancreatic secretory parenchyma and results in the replacement with the fibrous cells. This condition leads to the Diabetes mellitus [87]. Pancreatitis starts as the pancreatitis and cause the inflammation. On the other hand, reactive oxygen species (ROS) are the triggering factors for the pancreatitis, they are the enhancers of inflammation that have the potential to convert the impaired acinar cell into a manufactory for the cytokines and chemokines [26]. Such type of the inflammatory responses are the instigating factors for the beta cell disruption and resultantly, low levels of insulin and hyperglycemic condition [27-30]. However, there is lack of systematic studies that are capable of examining the main genetic differences between type 3c diabetes secondary to chronic pancreatitis.



Pancreatic Ductal Adenocarcinoma and Type 3c Diabetes Mellitus

Pancreatic cancer is the leading cause of wide scale deaths around the world and its risk is high in the patients with the new-onset diabetes. Therefore, it is named as pancreatic ductal adenocarcinoma (PDAC)-

associated diabetes mellitus (PDAC-DM), a “type 3C diabetes”. The onset of “PDAC-DM” mostly occurs 2–5 years prior to the diagnosis of “PDAC” that’s why its early diagnosis is very significant. However, it is not possible to differentiate PDAC-DM from the other diabetes on the basis of clinical signs and symptoms

so, there is the urgent need of the specific markers in the health sector [31-35].

Cystic Fibrosis and Type3c Diabetes Mellitus

Cystic fibrosis diabetes is the most apparent form of diabetes in people with cystic fibrosis. Although it possesses characteristics of both type 1 and type 2, but it is a quite different condition. Type 3c diabetes develops mainly due to the damage of pancreas that can happen because of a few typical reasons. Although it is much different from other types, still you can get a wrong diagnosis of diabetes type 2 because Type 3c isn't well diagnosed. Becoming unable to get the right diagnosis can be really hard to deal with it emotionally. You may feel angry for not having the right treatment or maybe you could just get ruined by the whole process. Thus, make sure you find the right person to communicate with.

Clinical Presentation of Type 3c Diabetes Mellitus

Large no of patients suffering from the type 3c Diabetes mellitus have some history of pancreatitis along with abdominal pain, steatorrhea or maldigestion with nutritional deficiencies and glucose intolerance. Patients may also flex the symptoms of maldigestion or abdominal pain without any diagnosis of chronic pancreatitis, or even may be asymptomatic except for glucose intolerance, and only through careful clinical evaluation is pancreatic disease suspected [36-40].

Symptoms of Diabetes Type 3c

Diabetes type 3c is a complex situation in which this is very difficult to diagnose that actually what the problem is. In specific, there are a number of symptoms that are associated with PEI (Pancreatic exocrine insufficiency). In this disease, the pancreas is not functional anymore and it does not provide the body with essential enzyme required for appropriate digestion. This is mainly due to chronic pancreatitis and cystic fibrosis. That is the reason why the symptoms of Diabetes type 3c are usually linked with digestive tract symptoms. The major symptoms include [41]:

- Diarrhea
- Abdominal bloating
- Eating issues
- Bloating and abdominal pain
- Nausea
- General fatigue

These symptoms are usually accompanied by a number of conditions like the patients with a history

pancreatic disorder, an instance of weight loss and a severe pain [42-45].

Diagnosis of Diabetes Type 3c

Diagnosis of Diabetes type 3c is too complicated and that's why it renders it undiagnosed and maltreated in most of the cases. A few reasons of its being difficult to diagnose is high variability of glucose level in the blood i.e., it sometimes becomes too high and sometimes become too low. That's why it sometimes sounds like Diabetes type 1 and hyperglycemia and sometimes it seems to be hyperglycemic. Now, after a long run of research there has been set a few standard steps to follow when diagnosing for Diabetes [46] which are:

1. Check for pancreatic exocrine insufficiency.
2. Perform pathological pancreatic imaging.
3. Check for Diabetes type 1-associated auto-antibodies.

Based on these criteria there can be a number of ways to check for type 3c and rule out the other types, for example absence of pancreatic enzymes in body after a glucose or mixed meal can be a true sign of type 3c. [47] HbA1C was initially thought to be a screening test for DM type 3c but there is a much lower connection between A1c and insulin tolerance levels, that's why HbA1c can't be used as a standard technique to diagnose DM type 3c [48]. It is simply conclusive that, the major relation of diabetes type 3c is with PEI and pancreatic disorders, so diagnosing in depth for the problems of associated with pancreatic problems is the major gateway to diagnose type 3c [74].

General Overview of The Differential Criteria Between T2 and T3c DM

Type 3c is characterized by deficiency of PP (pancreatic polypeptide) whereas type 2 contains high levels of nutrient stimulated pancreatic polypeptide i.e., PP. There are different criteria on which Type 2 DM and Type 3c DM can be categorized, which includes the pathological imaging i.e., endoscopy, MRI and ultrasound etc. Absence of some autoimmune markers associated with Type 1 DM is another basis of differentiation between these two. Some minor criteria include the identification of B-cell functionality, IR (Insulin Resistance), Abnormalities of incretin secretion and low levels of fat-soluble vitamins (A, D, E, K). However, Type 3c and Type 2 DM still overlaps at many points, which will need more research work to further differentiate the different types of diabetes [49-55].


Management of Diabetes Type 3c

Management of Diabetes type 3c is not a much-researched field due to less reported and diagnosed cases [56]. Management is based upon the cause of particular type 3c diabetes. Usually, it is suggested for the patients to get some knowledge and awareness about the diabetes management in sessions about the diabetes awareness programs and seminars [57-60]. Primarily the basic need is to maintain a little greater than the normal level of glucose in body i.e., to avoid hypoglycemia [78]. On the other hand, it is suggested for the patients with a history of chronic pancreatitis to take a diet that is rich in dietary fiber and lacks fat. In this way they can avoid the symptoms of steatorrhea, and they must also need to prevent hyperglycemia through diet [61]. Thus, it can be said that the only way to manage this least researched form of diabetes is to use a multi-dimensional approach in which both nutritional and therapeutic parameters

are taken under use [62]. And to minimize the serious complications, strategic management should be balanced to maintain the maximum glycemic control. Other healthy improvisations can also be helpful to prevent hyperglycemia [63-70].

Treatment of Diabetes Type 3c

About 75% of the type 3c diabetes result from the chronic pancreatitis which ultimately increases the risk of development of carcinoma in the pancreas. Treatment with insulin or insulin secretagogue can readily increase the risk of malignancy due to their hypoglycemic activity, whereas metformin could be beneficial in reducing it. However, in advanced type 3c DM insulin replacement therapy is preferred, which helps the patients achieve optimal glucose concentration in blood by mimicking the physiological delivery of insulin in a very comprehensive approach [71-81].



Conclusion

Diabetes mellitus is a particularly heterogeneous and multifactorial metabolic disorder that is characterized by a combination of insulin insufficiency and insulin resistance. The prevalence of diabetes mellitus has risen exponentially over the last three decades, with resultant increase in morbidity and mortality mainly due to its complications. There is need of Increases awareness of these complications among patients and populations, this facilitates early detection and prevention of these diseases. The earlier diagnosis, timely appropriate treatment may help in avoiding severe, irreversible disability and deformity, which can be done with the tool of patient education and conducting seminar to increase awareness.

References

1. Suhail, M., Riaz, S. (2020). In Silico Proteomics EVOO Therapy for Lipid Lowering in the Patients of Diabetes Mellitus. In *Biochemical Testing-Clinical Correlation and Diagnosis*. IntechOpen.
2. Johnson, C. D., Arbuckle, R., Bonner, N., Connett, G., Dominguez-Munoz, E., et al. (2017). Qualitative Assessment of The Symptoms and Impact of Pancreatic Exocrine Insufficiency (PEI) To Inform the Development of a Patient-Reported Outcome (PRO) Instrument. *The Patient-Patient-Centered Outcomes Research*, 10:615-628.
3. Conlon, K. C., Duggan, S. N. (2017). Pancreatogenic Type 3c Diabetes:

- Underestimated, Underappreciated and Poorly Managed. *Practical Gastroenterol*, 15.
4. Ewald, N., Kaufmann, C., Raspe, A., Kloer, H. U., Bretzel, R. G., et al. (2012). Prevalence of Diabetes Mellitus Secondary to Pancreatic Diseases (Type 3c). *Diabetes/Metabolism Research and Reviews*, 28(4):338-342.
 5. Ewald, N., Kaufmann, C., Raspe, A., Kloer, H. U., Bretzel, R. G., et al. (2007). Prevalence of Diabetes Mellitus Secondary to Pancreatic Diseases (Type 3c). *Experimental and Clinical Endocrinology & Diabetes*, 115(S 1):P01_028.
 6. Moran, A., Brunzell, C., Cohen, R. C., Katz, M., Marshall, B. C., et al. (2010). Clinical Care Guidelines for Cystic Fibrosis-Related Diabetes: A Position Statement of The American Diabetes Association and A Clinical Practice Guideline of The Cystic Fibrosis Foundation, Endorsed by The Pediatric Endocrine Society. *Diabetes Care*, 33(12):2697-2708.
 7. Rickels, M. R., Bellin, M., Toledo, F. G., Robertson, R. P., Andersen, D. K., et al. (2013). Detection, Evaluation and Treatment of Diabetes Mellitus in Chronic Pancreatitis: Recommendations from Pancreas Fest 2012. *Pancreatology*, 13(4):336-342.
 8. Makuc, J. (2016). Management of Pancreatogenic Diabetes: Challenges and Solutions. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 311-315.
 9. Andersen, D. K., Andren-Sandberg, Å., Duell, E. J., Goggins, M., Korc, M., et al. (2013). Pancreatitis-Diabetes-Pancreatic Cancer: Summary of an NIDDKNCI Workshop. *Pancreas*, 42(8):1227-1237.
 10. Golden, S. H., Robinson, K. A., Saldanha, I., Anton, B., Ladenson, P. W. (2009). Prevalence and Incidence of Endocrine and Metabolic Disorders in The United States: A Comprehensive Review. *The Journal of Clinical Endocrinology & Metabolism*, 94(6):1853-1878.
 11. Forouhi, N. G., Wareham, N. J. (2010). Epidemiology of Diabetes. *Medicine*, 38(11):602-606.
 12. American Diabetes Association. (2005). Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 28(Suppl 1):S37-4S2.
 13. Liu, Q. C., Zhuang, Z. H., Zeng, K., Cheng, Z. J., Gao, F., et al. (2009). Prevalence of Pancreatic Diabetes in Patients Carrying Mutations or Polymorphisms of The PRSS1 Gene in The Han Population. *Diabetes Technology & Therapeutics*, 11(12):799-804.
 14. Hart, P. A., Bellin, M. D., Andersen, D. K., Bradley, D., Cruz-Monserrate, Z., et al. (2016). Type 3c (Pancreatogenic) Diabetes Mellitus Secondary to Chronic Pancreatitis and Pancreatic Cancer. *The lancet Gastroenterology & Hepatology*, 1(3):226-237.
 15. Domschke, S., Stock, K. P., Pichl, J., Schneider, M. U., Domschke, W. (1985). Beta-Cell Reserve Capacity in Chronic Pancreatitis. *Hepato-Gastroenterology*, 32(1):27-30.
 16. Ewald, N., Kaufmann, C., Raspe, A., Kloer, H. U., Bretzel, R. G., et al. (2012). Prevalence of Diabetes Mellitus Secondary to Pancreatic Diseases (Type 3c). *Diabetes/Metabolism Research and Reviews*, 28(4):338-342.
 17. Sarles, H. (1986). Etiopathogenesis and Definition of Chronic Pancreatitis. *Digestive Diseases and Sciences*, 31(9):91-107.
 18. Braganza, J. M., Lee, S. H., McCloy, R. F., McMahon, M. J. (2011). Chronic Pancreatitis. *The Lancet*, 377(9772):1184-1197.
 19. Sasikala, M., Talukdar, R., Radhika, G., Rao, G. V., Pradeep, R., et al. (2012). β -Cell Dysfunction in Chronic Pancreatitis. *Digestive Diseases and Sciences*, 57(7):1764-1772.
 20. Gao, W., Zhou, Y., Li, Q., Zhou, Q., Tan, L., et al. (2015). Analysis of Global Gene Expression Profiles Suggests a Role of Acute Inflammation in Type 3C Diabetes Mellitus Caused by Pancreatic Ductal Adenocarcinoma. *Diabetologia*, 58(4):835-844.
 21. Gudipaty, L., Rickels, M. R. (2015). Pancreatogenic (Type 3c) Diabetes. *Pancreapedia: The Exocrine Pancreas Knowledge Base*.
 22. Riaz, S. And Mughal, M.S. (2000). Histopathological Changes in Kidney of Grass Carp, *Ctenopharyngodon Idella* as Manifested by Chromium Intoxication. *Biologia*, 46(1&2):83-88.
 23. Riaz, S. And Mughal, M.S. (2002). Histopathological Changes in Spleen of Grass Carp, *Ctenopharyngodon Idella* as Manifested by Chromium Intoxication. *Biologia*, 48(1&2):401-405.
 24. Riaz, S., Mughal, M.S. (2002). Effect of Different Levels of Tannery Wastes on Serum Enzymes of Quail Chicks *Coturnix Japonica*. *Biologia*, 2002. 48(1&2):389-400.

25. Riaz, S., Mughal, M.S. (2002). Histopathological Changes of Different Levels of Tannery Wastes Replaced by Chrome Shavings in Liver & Kidney of Quail Chicks *Coturnix Japonica*. *Biologia*, 48(1&2):341-349.
26. Riaz, S., Riaz, A. (2006). Effect of Different Levels of Chrome Shaving on Total Body Weight of Quail Chicks *Coturnix Japonica*. *Journal of Biochemistry and Molecular Biology*. 39(1&2):10-19.
27. Riaz, S., Shoukat S., Mughal, M.S. (2007). Fluoride Toxicity on The Histopathological Aspects of Liver in The Cirrhinus Mrigala. *Journal of Biochemistry and Molecular Biology*. 39(3):165-173.
28. Riaz, S., Shoukat S., Mughal, M.S. (2008). Fluoride Toxicity on The Histopathological Aspects of Kidney in The Cirrhinus Mrigala. *Journal of Biochemistry and Molecular Biology*. 40(3):20-24.
29. Riaz, S., Alam S.S., Rabbani N., Thornalley P.J., Aktar M.W. (2009). Proteomic Strategies to Study the Protein Biomarker for Diabetes Mellitus and Effect of Thiamine on Their Levels. *Proceeding of Symposium of Genomics, Proteomics and Metabolomics. Modern Trends in Biotechnology. Dept. Of MMG, University of The Punjab, Lahore*. 198-212.
30. Riaz, S., Alam, S.S., Ikram, A. (2006). Histopathological Changes Observed in The Heart and Gizzard of Quail Chicks *Coturnix Japonica* Administrated by The Different Levels of Chrome Shaving. *African Journal of Biotechnology*. 5(19):1765-1769.
31. Riaz, S., Alam, S.S. (2006). Evaluation of Nutritive Value of Different Levels of Chrome Shaving Partially Replacing Animal Protein in The Feed on The Performance of Quail Chicks. *International Journal of Agricultural and Biological Science*, 1(1):1-7.
32. Riaz, S., Alam, S.S., Ali, G. (2006). Effect of Different Levels of Chrome Shavings on Hematological Parameters in The Blood of Quail Chicks. *International Journal of Agricultural and Biological Science*, 1(1):27-32.
33. Naila Rabbani, Saadia Shahzad Alam, Samreen Riaz, James Larkin, M Waheed Akhtar, et aly. (2009). High Dose Thiamine Therapy for People with Type 2 Diabetes and Microalbuminuria: A Randomised, Double-Blind, Placebo-Controlled Study. *Diabetologia. Springer Berlin / Heidelberg*. 52(2):208-212.
34. Samreen Riaz, Mehreen Raza, Saadia Shhazad Alam, Shahida Hasnain, M. Waheed Akhtar. (2009). Obesity as Risk Factor and Study of Obesity Related Proteins in Diabetes Mellitus. *African Journal of Biotechnology*. 8(5):737-744.
35. Nawaz, K., Riaz, S., Hasnain, S. (2009). Screening of Anti Methicillin Resistant *Staphylococcus Aureus* (MRSA) Bacteriocin Producing Bacteria. *African Journal of Biotechnology*. 8(3):365-368.
36. Rabbani, N., Shahzad Alam, S., Riaz, S., Larkin, J.R., Akhtar, M.W., et al. (2009) High Dose Thiamine Therapy for Patients with Type 2 Diabetes and Microalbuminuria: A Pilot Randomised, Double-Blind, Placebo-Controlled Study. *Diabetologia*, 52(2):208-212.
37. Harman, C. (2009). Could Vitamin 'B' the Answer? *Nature Reviews Endocrinology*, 5(5):236-236.
38. Harman, C. (2009). Could Vitamin 'B' the Answer? *Nature Reviews Nephrology*. 5(4):182.
39. Riaz, S. (2009). Review of Diabetes Mellitus. *Academic Journals*, 4(5):367-373.
40. Samreen Riaz, Saadia Shahzad Alam, M. Waheed Akhtar. (2010). Proteomic Identification of Human Serum Biomarkers in Diabetes Mellitus Type-2. *Journal of Pharmaceutical and Biomedical Analysis*. 51(5):1103-1107.
41. Samreen Riaz, Ansa Butt. (2010). Study of Protein Profiling of Human Urine in Diabetic Hypertensive Nephropathy Versus Normal Healthy Controls. *Diabetes Technology and Therapeutics*. 12(5):379-386.
42. Samreen Riaz, Saadia Shahzad Alam, Surjit Kaila Srai, Vernon Skinner, Aasma Riaz et al. (2010). Proteomic Identification of Human Urine Biomarkers in Diabetes Mellitus Type-2. *Journal of Diabetes Technology and Therapeutics*. 12(12):979-988.
43. Samreen Riaz, Vernon Skinner, Surjit Kaila Srai. (2011). Effect of High Dose Thiamine on Levels of Human Urine Protein Biomarkers in Diabetes Mellitus Type 2. *Journal of Pharmaceutical and Biomedical Analysis*. 54(4):817-825.
44. Samreen Riaz, Vernon Skinner, Surjit Kaila Srai. (2011). Corrigendum to "Effect of High Dose Thiamine on The Levels of Urinary Protein Biomarkers in Diabetes Mellitus Type 2", *Journal of Pharmaceutical and Biomedical Analysis*. 56(1):139.

45. Saadia Shahzad Alam, Samreen Riaz, M. Waheed Akhtar. (2011). Effect of High Dose Thiamine Therapy on Activity and Molecular Aspects of Transketolase in Type 2 Diabetic Patients. *African Journal of Biotechnology*. 10(75):17305-17316.
46. Saadia Shahzad Alam, Samreen Riaz, M. Waheed Akhtar. (2012). Effect of High Dose Thiamine Therapy on Risk Factors in Type 2 Diabetics. *Journal of Diabetes and Metabolism*, 3:10.
47. Saadia Shahzad Alam, Samreen Riaz, M. Waheed Akhtar. (2013). Effect of High Dose Thiamine Therapy on Induction and Activities of Pyruvate Dehydrogenase and A-Ketoglutarate Dehydrogenase in Type 1 Diabetic Patients. *British Journal of Medicine & Medical Research*. 12(10):1-12.
48. Saadia Shahzad Alam, Samreen Riaz, M. Waheed Akhtar. (2013). Effect of High Dose Thiamine Therapy on Activity and Molecular Aspects of Transketolase in Type 1 Diabetic Patients. *African Journal of Biotechnology*. 12(75):17305-17316.
49. Saadia Shahzad Alam, Samreen Riaz, M. Waheed Akhtar. (2013). Effect of High Dose Thiamine Therapy on Induction and Activities of Pyruvate Dehydrogenase and A-Ketoglutarate Dehydrogenase in Type 2 Diabetic Patients. *Diabetes Technology & Therapeutics*. 12(12):979-988.
50. Saadia Shahzad Alam, Samreen Riaz, M. Waheed Akhtar. (2013) Effect of High Dose Thiamine Therapy on Induction and Activities of Pyruvate Dehydrogenase and A-Ketoglutarate Dehydrogenase in Type 1 Diabetic Patients. *Diabetes Educator*. 12(1):1-15.
51. Samreen Riaz, Sadia Rana. (2014). Analysis of Uropathogens Among Type II Diabetic Patients in Pakistani Population. *Diabetes And Metabolism*, 2(2):31-41.
52. Samreen Riaz. (2014). Obesity As a Risk Factor for Diabetes Mellitus in The Local Population of Pakistan. *Universal Journal of Clinical Medicine*. 2(3):58-64.
53. Riaz S. (2014). Evaluation and Analysis of Human Folate Levels in Pakistani Diabetic Population. *International Journal of Scientific & Engineering Research*, 5(12):1572-1576.
54. Riaz S. (2015), Study of Protein Biomarkers of Diabetes Mellitus Type 2 and Therapy with Vitamin B1. *Journal of Diabetes Research*. 1-10.
55. Saadia Shahzad Alam, Samreen Riaz (2016). Induction and Activities of Pyruvate Dehydrogenase And A-Ketoglutarate Dehydrogenase in Type 2 Diabetic Patients and Therapy with Vitamin B1. *International Journal of Advances in Medicine and Medical Research*. 12(10):1-10.
56. Mursleen, M. T., Riaz, S. (2017). Implication of Homocysteine in Diabetes and Impact of Folate and Vitamin B12 in Diabetic Population. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 11:S141-S146.
57. Riaz S. (2018). Therapeutic Implication of Folate-Homocysteine Interaction in The Local Diabetic Pakistani Population. *Annals of Vascular Medicine Research*. 5(1):1085-1089.
58. Afifa, Samreen Riaz. (2018). Analysis Of Pyridoxine in The Male Diabetic Population of Lahore and Sheikhpura. *Annals of Diabetes Metabolic Disorders & Control*. 2(1):118.
59. Riaz S., Tariq M. (2018). Linkage of Micro Albuminuria and Serum Albumin Levels in The Diabetic Patients Punjab University Premises. *Annals of Diabetes Metabolic Disorders & Control*. 2(1):117.
60. Riaz S., Tariq M., Aslam S. Association of Serum Protein Levels in The Diabetic Patients with Risk of Cardiovascular Disease and Nephropathy in Pakistani Population. *Diabetes And Metabolism Journal*. 4(1):11-15.
61. Afifa, Samreen Riaz. (2018). Analysis of Pyridoxine in The Male Diabetic Population of Lahore and Sheikhpura. *Annals of Diabetes Metabolic Disorders & Control*. 2(1):118.
62. Haseeba Yousaf, Samreen Riaz. (2018). Biomolecular Evaluation of Cobalamin (Vitamin B12) In the Diabetic Population of Lahore and Sheikhpura. *Annals of Diabetes Metabolic Disorders & Control*. 2(1):119.
63. Samreen Riaz, Shagufta Naz, Umera Kousar, Zil E Huma Malik, Rasheeda Bashir. (2018). Major Complication Associated with Diabetes Mellitus Type II in Punjab and Khyber Paktunkhwa Population. *Diabetes and Metabolism*, 9:809.
64. Javaria Yasin, Samreen Raiz. (2019). Evolutionary History and Molecular Epidemiology of HIV Pol Gene in Pakistan. *Pakistan Journal of Medical Sciences*. 15(1):1-10.
65. Samreen Riaz, Amina Mehboob. (2020). Olive Oil Tends to Lowers Cholesterol Level in Patients Suffering from Diabetes. *Science Alert. International Journal of Biological Sciences*. 1-10.

66. Nusair Malik, Arwa Faisal, Snobia Munir, Samreen Riaz. (2020). Clinical and Microbiological Profile of Urinary Tract Infection in Male Diabetic Patients. *Journal of Microbiology and Experimentation*. 8(1):30-34.
67. Farah Qudsia, Hafiza Amina Matti, Samreen Riaz. (2020). Pleiotropic Effect of Folate-Cobalamin Combinational Therapy on Diabetes Mellitus. *Annals of Nutrition Disorders and Therapy*. 7(1):1061.
68. Snobia Munir, Samreen Riaz. (2020). Type 2 Diabetes and Cancer. *Annals of Clinical Endocrinology and Metabolism*. 4:1-6.
69. Snobia Munir, Samreen Riaz. (2020). Molecular Association of Diabetes Mellitus with Different Types of Cancer from Population of Lahore. *Annals Of Diabetes, Metabolic Disorders and Control*. 3(1):126.
70. Hafiza Amina Matti, Samreen Riaz. (2022). Molecular Association of Cancer and Diabetes Mellitus with Combinational Therapy. *International European Extended Enablement-In Science, Engineering & Management*, 8(4).
71. Hafiza Amina Matti, Samreen Riaz. (2022). Evaluation of Environmental Influence of Arsenic Compounds on Patients Suffering from Type 2 Diabetes. *International Journal of Scientific & Engineering Research*.
72. Samreen Riaz, Shagufta Naz, Umera Kousar, Zil E Huma Malik, Rasheeda Bashir. (2020). The Association Between VEGF Rs699947 Polymorphism and Their Serum Level in Patients with Diabetic Foot Ulcer in Pakistani Population.
73. Samreen Riaz, Abdullah Mohsin, (2020). Role of Phytochemicals Derived from Plants Biodiversity and Its Impact in Treating COVID-19. *Journal of Bioscience & Biomedical Engineering*. 1(3)1-8.
74. Nusair Malik, M. S., Zubair, M., Riaz, S. (2020). Therapeutic Interventions of EVOO As Nutraceutical Agent to Lower Lipid Profile in The Diabetes Mellitus Type 2. *Journal of Bioscience & Biengineering*, 1(3):1-8.
75. Munir S, Riaz S, Arshad T, Riaz A. (2020). Risk Factors Associated to Patients with Type 2 Diabetes In Lahore District. *Annals of Clinical Endocrinology and Metabolism*. 4:11-19.
76. Samreen Riaz. (2020). Molecular Association of Coronavirus with Diabetes Mellitus Type 2 In Pakistan *Journal of Diabetes Research*. 2(4):1-11.
77. Areeba Saeed, Sohaib Imtiaz, Samreen Riaz. (2020). Study of Urinary Tract Infection in Female Patients Suffering from Cancer. *Journal of Cancer Research Reviews & Reports*. 2(2):1-15.
78. Sohaib Imtiaz, Samreen Riaz. (2020). Antimicrobial Susceptibility Profile of Bacteria Isolated from Male Cancer Patients. *International Journal of Clinical Studies & Medical Case Reports*. 6(3):1-6.
79. Riaz, S., ul Arsalan, S., Ali, R., Riaz, A. (2020). Statistical Analysis to Identify the Effect of Risk Factors on Diabetic Patients from the Sheikh Zayed Hospital Lahore. *Current Trends on Biostatistics and Biometrics*, 3(3):368-382.
80. Iqbal, R., Aslam, S., Liaqat, I., Zafar, A., Aftab, K., & Riaz, S. (2021). Polymorphisms in CDKN2B-AS1 gene Confer a Risk for Myocardial Infarction in Pakistani Population. *Bioscience Reports*. BSR, 2003.
81. Shokat, S., Iqbal, R., Yaqub, A., Shahid, F., Riaz, S. (2021). Analysis of Environmental Pollutants as Diabetogenic Agent in The Patient Suffering from Diabetes Mellitus. *Diabetes & Obesity International Journal*, 6(1).

Cite this article: Riaz S. (2025). Prevalence and Epidemiology of Diabetes Mellitus Type 3C, *International Journal of Biomedical and Clinical Research*, BioRes Scientia Publishers. 3(3):1-10. DOI: 10.59657/2997-6103.brs.25.052

Copyright: © 2025 Samreen Riaz, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Article History: Received: January 09, 2025 | Accepted: January 31, 2025 | Published: February 07, 2025